

BOSE-FERMI SYMMETRY: A CRUCIAL ELEMENT IN ACHIEVING UNIFICATION[†]

Jogesh C. Pati

Institute for Advanced Study
Princeton, NJ 08540 USA
and

Department of Physics, University of Maryland*
College Park, MD 20742 USA

ABSTRACT

This talk is dedicated to honor the memory of Professor S.N. Bose. I survey the crucial roles played by Bose-Fermi symmetry in all recent attempts at higher unification, which include the ideas of (i) the conventional approach to grand unification; (ii) the preonic approach; and (iii) superstrings.

[†] Talk presented at the Int'l. Sym. on "Bose and Twentieth Century Physics," Calcutta, India, January 1994, to appear in the Proceedings.

* Permanent address. E-mail: "pati@umdhep.umd.edu"

I. Preliminary Remarks

I regard it a special privilege to speak at this conference which is dedicated to celebrate the birth centenary of a great scientist and my country-man – Professor Saytendra Nath Bose. Beyond doubt, Bose’s contribution [1] to physics is one of the landmarks in the development of quantum theory. In one stroke, he introduced two new concepts of lasting value: the concept of massless particles with two states of polarization (these are the photons with spin 1) and the concept that their number is not conserved; that they obey a new statistics. In Pais’ words, “The paper by Bose is the fourth and last of the revolutionary papers of the old quantum theory (the other three being by respectively Planck, Einstein and Bohr)” [2]. To this I will add that Bose’s contribution, attributing the concept of a new statistics to photons, turned out to be an integral feature of relativistic quantum field theory as well through a realization that evolved during the 1930’s through the 50’s. This is the famous connection between spin and statistics [3] which asserts that particles of integer spins (i.e, spin 0,1,2,...) obey what is now called the Bose-Einstein statistics while those of half-integer spins (i.e.,spin-1/2, 3/2, 5/2, ...) obey the Fermi-Dirac statistics or the exclusion principle. This in turn allows us to classify all known elementary particles as either *bosons* or *fermions* which respectively obey the Bose-Einstein and the Fermi-Dirac statistics.

In honor of Professor Bose, I will elucidate here the role of a further concept in recent attempts at achieving a unification of matter and its forces. This is the concept of a symmetry that relates fermions to bosons, which evolved some fifty years after the birth of quantum statistics.

As a prelude, let me first say a few words about the status of the field of particle physics as it existed before the introduction of this new symmetry. One central goal of elementary particle physics has been to search for principles which would dictate the existence of particles and their forces. To cite a few examples, the principle of local gauge invariance dictates the existence of the photon and the gluons as well as that of the associated forces of quantum electro- and chromodynamics. The

principle of general coordinate invariance proposed by Einstein leads to the familiar gravitational “forces” as a consequence of the curvature of space-time and dictates the existence of spin-2 graviton. These principles in turn help preserve the masslessness of these “gauge” particles despite quantum corrections, at least in perturbation theory. Coming now to spin-1/2 particles, although there was no such *a priori* rationale, at least not until the developments in the 70’s and the 80’s for their existence, one may advance a different type of reason which is that spin-1/2 is the *smallest* unit of spin, associated with an elementary particle, that one needs to build all higher spin-particles as composites. Furthermore, purely from an utilitarian point of view, spin-1/2 particles (i.e., electrons) are at least *needed* since they obey the exclusion principle which is relevant to the explanation of chemistry and in turn to the biology of life. Now, once spin-1/2 particles are introduced into the lagrangian, they have the good feature that their masses remain protected against arbitrarily large quantum corrections through the so-called chiral symmetry which guarantees that the quantum corrections to the masses of spin-1/2 fermions in perturbation theory either vanish or are bounded by a logarithmic cutoff (symbolizing short-distance physics) depending upon whether their “bare” mass-terms are zero or non-zero.

By contrast, no such principle, not even utilitarian arguments, existed until the early 70’s, which would either dictate the existence of elementary spin-0 bosons or guard their masses against large quantum corrections. Nevertheless, there, of course, exist spin-0 particles which are known to be relevant in particle physics. In particular, the pions are the carriers of the nuclear force and the Higgs bosons (yet to be discovered) induce spontaneous breaking of the electroweak symmetry and give masses to W and Z bosons. As regards the pions, it is known that they are *not* elementary. They can be identified as the (pseudo) goldstone bosons associated with a dynamical breaking of the chiral symmetry of the up and down quarks and thus can be viewed as $q\bar{q}$ composites. For this reason, the mass of the pion does not get large quantum corrections. It is controlled by the relevant chiral symmetry breaking parameter which is determined by (a) the finite “bare” masses of the up and down

quarks and (b) the QCD scale-parameter.

What about the origin of the Higgs-boson? It is not yet known (especially if one allows for the preonic idea to be mentioned shortly) whether the Higgs boson is elementary or composite. The idea of compositeness of the Higgs boson in the sense of technicolor, which treats quarks, leptons and technifermions as elementary but Higgs boson as composite is perhaps excluded because it runs into difficulties with flavor-changing neutral current processes and oblique electroweak parameters. As I shall elucidate later in this talk, no such difficulty exists, however, and one obtains a viable and economical picture, if one assumes that, together with the Higgs boson, the quarks and the leptons are composite as well, sharing common constituents – called “preons”. Thus, either the Higgs boson is composite in the context of a certain preonic theory, or it is elementary. In case it is elementary, which is in fact the conventional view, the two pertinent questions are: (i) What if any is an *a priori* rationale for its existence, and, equally important, (ii) how can one protect its mass against large quantum corrections? To be specific, since the Higgs boson couples to the gauge bosons and also possesses quartic self couplings, one obtains corrections in one loop to its mass which are proportional to $\alpha_i \Lambda^2$ (Λ is the cutoff characterizing short-distance physics and α ’s are coupling parameters). Allowing for short-distance physics to include at least gravity and possibly grand unification, one would expect $\Lambda \sim 10^{16} - 10^{19} \text{ GeV}$. To obtain a physical mass $\lesssim 1 \text{ TeV}$, one thus needs *unnatural fine tuning* by some 24 orders of magnitude (or higher) in the choice of the bare mass of the Higgs boson to cancel the large quantum corrections. Such a fine tuning is unnatural, unattractive and thus unacceptable in a fundamental theory.

The problem of this unnatural fine tuning gets resolved and the existence of spin-0 bosons derives a significance on a par with that of spin-1/2 fermions through the idea of a new symmetry – commonly called “supersymmetry” [4] – which transforms spin-1/2 fermions into spin-0 bosons and *vice versa*. Since it is a symmetry that transforms bosons into fermions, I will sometimes refer to it in the course of this talk as the “*Bose-Fermi Symmetry*”.

The power of supersymmetry arises because its generator(s) Q transforms a spin-0 (or spin-1) boson into a spin-1/2 fermion and thereby changes the spin of the particle by 1/2 unit as well as its statistics. Thus, Q transforms as a spin-1/2 fermionic operator. This is in contrast to the generators of the time-honored Lie algebras, associated with the familiar symmetries such as isospin and $SU(3)$, whose generators transform as Lorentz-scalars – i.e., as spin-0 bosonic operators; and which can thus transform a particle of a given spin into another of the *same spin*, only. The fermionic generators $(Q, \bar{Q})$ of $N = 1$ supersymmetry and the bosonic energy-momentum operators P_μ , together, define in fact a graded Lie algebra, consisting of a combination of commutators and anticommutators. In particular, they satisfy: $\{Q, \bar{Q}\} = 2\sigma^\mu P_\mu$; $\{Q, Q\} = \{\bar{Q}, \bar{Q}\} = 0$; $[P_\mu, Q] = [P_\mu, \bar{Q}] = [P_\mu, P_\nu] = 0$.

Because of this new feature, supersymmetry brings some major benefits:

- (i) First, as mentioned above, supersymmetry unites fermions and bosons as members of a supermultiplet and thereby provides the rationale for the existence of spin-0 matter, *on a par* with that of spin-1/2 matter.
- (ii) Second, supersymmetry permits a non-trivial marriage of space-time (Poincaré) symmetries with internal symmetries in accord with relativity. As shown by Haag, Lopuszanski and Sohnius [5], the graded Lie algebra associated with supersymmetry is the only framework within which such a marriage can be achieved consistent with relativistic quantum field theory. For example, $SU(2)$ -isospin symmetry together with supersymmetry groups the spin-1/2 doublet of (u, d) -quarks with the spin-0 doublet of $(\tilde{u}, \tilde{d})$ -squarks to make a super-bidoublet

$$\begin{bmatrix} u & \leftrightarrow & \tilde{u} \\ \updownarrow & & \updownarrow \\ d & \leftrightarrow & \tilde{d} \end{bmatrix}$$

where all four members are degenerate and are related to each other by symmetry generators. To judge the importance of this property of supersymmetry, it is useful to recall past attempts of this nature which proposed to combine particles such as $(\pi$ and $\rho)$ and $(N$ and $N^*)$ which differ in spin only by integer units. These attempts failed because, as shown by Coleman and Mandula [6], with only bosonic symmetry

operators satisfying Lie algebras, they did not satisfy the constraints of relativity. In short, Bose-Fermi symmetry, together with the associated graded Lie algebra, brings about a synthesis of fundamental matter that goes well beyond that permissible within symmetries of just bosonic operators.

(iii) The third major advantage of quantum field theories with supersymmetry is that as a rule these theories are far less ultraviolet divergent than their non-supersymmetric counterparts. This comes about due to cancellation between fermionic and bosonic partners in quantum loops in SUSY theories [7]. Because of such cancellation, the $N = 4$ supersymmetric Yang-Mills theories turn out in fact to be ultraviolet finite, and point-particle supergravity theories, though still non-renormalizable, exhibit much better ultraviolet behavior than non-supersymmetric Einstein gravity. Owing to the same cancellation in SUSY theories, the quadratic divergence in the spin-0 boson self-energy (mass)² drops out. Instead, the self-(mass)² of spin-0 bosons is given by $(\alpha_i/\pi)|m_B^2 - m_F^2|$, where B and F respectively denote boson and fermion SUSY partners. This would, of course, vanish if SUSY were exact (i.e, $m_B = m_F$). In practice, it is controlled by the SUSY-breaking parameter $|m_B^2 - m_F^2|$. Assuming that this parameter is of order 1 to 10 TeV², say, we see that in SUSY theories one no longer needs unnatural fine tuning of the Higgs (mass)² to keep the mass of the Higgs-boson at the 1 TeV-scale. This feature is thus generally regarded as one of the main *practical motivations* for supersymmetry and the reason for expecting that SUSY partners should exist at the TeV-scale.

(iv) The fourth major advantage of supersymmetry – one that would be most needed if the idea of preonic substructure turns out to be right (see discussions later) – is certain novel properties of a class of strongly interacting supersymmetric gauge theories relative to non-supersymmetric QCD. To cite one such feature, the Witten index theorem forbids a dynamical breaking of supersymmetry in SUSY SU(N)-theories with massless matter, at least in so far as one can neglect gravity [8]. As a result, the matter-fermion condensate $\langle \bar{\psi}\psi \rangle$, which breaks not only chiral symmetry but also supersymmetry, must vanish in such theories (at least as $M_{Planck} \rightarrow \infty$). This is in

striking contrast to the case of ordinary QCD where the chiral symmetry-breaking quark-pair condensate $\langle \bar{q}q \rangle$ does in fact form with a normal strength. It is this inhibition in the formation of $\langle \bar{\psi}\psi \rangle$ and other SUSY-breaking condensates which has recently been utilized to build a viable and economical preon-model with many attractive features [9,10,11,12]. These include: (i) an explanation of the protection of composite quark-lepton masses compared to their scale of compositeness [10] and (ii) a natural origin of the hierarchy of mass-scales from $M_{Pl} \rightarrow m_W \rightarrow m_e \rightarrow m_\nu$ [9].

(v) Last but not least, the greatest benefits of Bose-Fermi symmetry is derived in the context of all attempts at higher unification, which include the ideas of (a) the conventional approach to grand unification, (b) the preonic approach, (c) supergravity and, of course, (d) superstrings. To put in one sentence, it seems that none of these ideas would work without the aid of supersymmetry. To present some of these benefits of supersymmetry in the context of higher unification, which is the main purpose of my talk, I need to say a few words about the puzzles in particle physics which confront us in the context of the standard model and the unifying ideas which have been proposed to resolve some of these puzzles.

II. Going Beyond the Standard Model

The standard model of particle physics (SM) has brought a good deal of synthesis in our understanding of the basic forces of nature, especially in comparison to its predecessors, and has turned out to be brilliantly successful in terms of its agreement with experiments. Yet, as recognized for some time [13], it falls short as a fundamental theory because it introduces some 19 parameters. And it does not explain (i) family replication; (ii) the coexistence of the two kinds of matter: quarks *and* leptons; (iii) the coexistence of the electroweak *and* the QCD forces with their hierarchical strengths $g_1 \ll g_2 \ll g_3$, as observed at low energies; (iv) quantization of electric charge; (v) inter and intrafamily mass-hierarchies - *i.e.*, $m_{u,d,e} \ll m_{c,s,\mu} \ll m_{t,b,\tau}$ and $m_b \ll m_t$, etc. - reflected by ratios such as $(m_u/m_t) \sim 10^{-4}$, $(m_c/m_t) \sim 10^{-2}$ and $(m_b/m_t) \sim \frac{1}{35}$; and (vi) the origin of diverse mass scales that span over more

than 27 orders of magnitude from M_{Planck} to m_W to m_e to m_ν , whose ratios involve very small numbers such as $(m_W/M_{Pl}) \sim 10^{-17}$, $(m_e/M_{Pl}) \sim 10^{-22}$ and $(m_\nu/M_{Pl}) < 10^{-27}$. There are in addition the two most basic questions: (vii) how does gravity fit into the whole scheme, especially in the context of a good quantum theory?, and (viii) why is the cosmological constant so small or zero?

These issues constitute at present some of the major puzzles of particle physics and provide motivations for contemplating new physics beyond the standard model which should shed light on them. The ideas which have been proposed and which do show promise to resolve at least some of these puzzles, include the following hypotheses:

(1) **Grand Unification:** The hypothesis of grand unification, which proposes an underlying unity of the fundamental particles and their forces [13,14,15], appears attractive because it explains at once (i) the quantization of electric charge, (ii) the existence of quarks *and* leptons with $Q_e = -Q_p$, and (iii) the existence of the strong, the electromagnetic and the weak forces with $g_3 \gg g_2 \gg g_1$ at low energies, but $g_3 = g_2 = g_1$ at high energies. These are among the puzzles listed above and grand unification resolves all three. *Therefore I believe that the central concept of grand unification is, very likely, a step in the right direction.* By itself, it does not address, however, the remaining puzzles listed above, including the issues of family replication and origin of mass-hierarchies.

(2) **Supersymmetry:** As mentioned before, this is the symmetry that relates fermions to bosons[4]. As a local symmetry, it is attractive because it implies the existence of gravity. It has the additional virtue that it helps maintain a large hierarchy in mass-ratios such as $(m_\phi/M_U) \sim 10^{-14}$ and $(m_\phi/M_{pl}) \sim 10^{-17}$, without the need for fine tuning, provided, however, such ratios are put in by hand. Thus it provides a technical resolution of the gauge hierarchy problem, *but by itself does not explain the origin of the large hierarchies.*

(3) **Compositeness:** Here there are *two distinct suggestions*:

(a) Technicolor: The idea of technicolor [16] proposes that the Higgs bosons are composite but quarks and leptons are still elementary. Despite the attractive feature of dynamical symmetry breaking which eliminates elementary Higgs bosons and thereby the arbitrary parameters which go with them, this idea is excluded, at least in its simpler versions, owing to conflicts with flavor-changing neutral current processes and oblique electroweak corrections. The so-called walking technicolor models may be arranged to avoid some of these conflicts at the expense, however, of excessive proliferation in elementary constituents. Furthermore, as a generic feature, none of these models seem capable of addressing any of the basic issues listed above, including those of family replication and fermion mass-hierarchies. Nor do they go well with the hypothesis of a unity of the basic forces.

(b) Preons: By contrast, the idea of preonic compositeness which proposes that not just the Higgs bosons but also the quarks and the leptons are composites of a *common* set of constituents called “preons” seems much more promising. Utilizing supersymmetry to its advantage, the preonic approach has evolved over the last few years to acquire a form [9,10,11,12] which is (a) far more economical in field-content and especially in parameters than either the technicolor or the conventional grand unification models, and, (b) is viable. Most important, utilizing primarily the symmetries of the theory (rather than detailed dynamics) and the peculiarities of SUSY QCD as regards forbiddenness of SUSY-breaking, in the absence of gravity, the preonic approach provides simple explanations for the desired protection of composite quark-lepton masses and at the same time for the origins of family-replication, inter-family mass-hierarchy and diverse mass scales. It also provides several testable predictions. In this sense, though still unconventional, the preonic approach shows promise in being able to address certain fundamental issues. I will return to it shortly.

(4) **Superstrings**: Last but not least, the idea of superstrings [17] proposes that the elementary entities are not truly pointlike but are extended string-like objects with sizes $\sim (M_{Planck})^{-1} \sim 10^{-33}$ cm. Strings with worldsheet supersymmetry, which constitute superstrings, possess considerable advantage over non-

supersymmetric strings in (a) stabilizing the vacuum and, (b) avoiding large quantum corrections to Higgs (mass)² through sub-Planck scale physics. Furthermore, the superstring theories automatically avoid tachyons. These theories (which may ultimately be just one) appear to be most promising in providing a unified theory of all the forces of nature including gravity and yielding a well-behaved quantum theory of gravity. In principle, assuming that quarks, leptons and Higgs bosons are elementary, a suitable superstring theory could also account for the origin of the three families and the Higgs bosons at the string unification scale, as well as explain all the parameters of the standard model. But in practice, this has not happened as yet. Some general stumbling blocks of string theories are associated with the problems of (i) a choice of the ground state (the vacuum) from among the many solutions and (ii) understanding supersymmetry breaking.

The ideas listed above are, of course, not mutually exclusive. In fact the superstring theories already comprise local supersymmetry and the central idea of grand unification. It remains to be seen, however, whether they give rise, in accord with the standard belief, to elementary quarks and leptons, or alternatively to a set of substructure fields – the preons. In the following, I first recall the status of conventional grand unification, and then provide a perspective as well as motivations for an alternative approach to grand unification, based on the idea of preons. In either case, I highlight the role of supersymmetry.

III. Grand Unification in the Conventional Approach and Supersymmetry

By “Conventional approach” to grand unification I mean the one in which quarks and leptons – and traditionally the Higgs bosons as well – are assumed to be elementary [13,14,15]. Within this approach, there are two distinct routes to higher unification: (i) the $SU(4)$ -color route [13] and (ii) $SU(5)$ [14]. Insisting on a compelling reason for charge – quantization, the former naturally introduces the left-right symmetric gauge structure $G_{224} = SU(2)_L \times SU(2)_R \times SU(4)_{L+R}^C$ [13], which in

turn may be embedded in anomaly-free simple groups like $SO(10)$ or E_6 [18].

It has been known for sometime that the dedicated proton decay searches at the IMB and the Kamiokande detectors [19], and more recently the precision measurements of the standard model coupling constants (in particular $\sin^2\hat{\theta}_W$) at LEP [20] put severe constraints on grand unification models without supersymmetry. Owing to such constraints, the non-SUSY minimal $SU(5)$ and, for similar reasons, the one-step breaking non-SUSY $SO(10)$ -model, as well, are now excluded beyond a shadow of doubt.

But the idea of the union of the coupling constants g_1, g_2 , and g_3 can well materialize in accord with the LEP data, if one invokes supersymmetry [21,22,23] into minimal $SU(5)$ or $SO(10)$. See Fig. 1, which shows the *impressive meeting* of the three coupling constants of the minimal supersymmetric standard model (MSSM) with an assumed SUSY-threshold around 1 TeV. Such a model can, of course, be embedded within a minimal SUSY $SU(5)$ or $SO(10)$ model, which would provide the rationale for the meeting of the coupling constants at a scale $M_U \approx 2 \times 10^{16}$ GeV, and for their staying together beyond that scale.

The fact that the coupling constants meet in the context of these models is reflected by the excellent agreement of their predicted value of $[\sin^2\hat{\theta}_W(m_z)_{theory} = .2325 \pm .005$ (using $\alpha_s(m_z) = .12 \pm .01$) with that determined at LEP: $[\sin^2\hat{\theta}_W(m_z)]_{expt.} = .2316 \pm .0003$. In SUSY $SU(5)$ or $SO(10)$, dimension 5 operators do in general pose problems for proton decay. But the relevant parameters of the SUSY-space can be arranged to avoid conflict with experiments [24]. The SUSY-extensions of $SU(5)$ or $SO(10)$ typically lead to prominent strange particle decay modes, e.g., $p \rightarrow \bar{\nu}K^+$ and $n \rightarrow \bar{\nu}K^0$, while a 2-step breaking of $SO(10)$ via the intermediate symmetry G_{224} can also lead to prominent $\Delta(B - L) = -2$ decay modes of the nucleon via Higgs exchanges such as $p \rightarrow e^-\pi^+\pi^+$ and $n \rightarrow e^-\pi^+$ and even $n \rightarrow e^-e^+\nu_e$, etc. in addition to the canonical $e^+\pi^0$ -mode [25].

It is encouraging that the super-Kamiokande (to be completed in April 1996) is expected to be sensitive to the $e^+\pi^0$ mode up to partial lifetimes of few $\times 10^{34}$

years, to the $\bar{\nu}K^+$ and $\bar{\nu}K^0$ modes with partial lifetimes $\leq 10^{34}$ years and to the non-canonical $n \rightarrow e^-e^+\nu_e$ and $p \rightarrow e^-\pi^+\pi^+$ modes with partial lifetimes $< 10^{33}$ years. Thus the super-Kamiokande, together with other forthcoming facilities, in particular, ICARUS, provide a *big ray of hope* that first of all one will be able to probe much deeper into neutrino physics in the near future and second proton-decay may even be discovered within the twentieth century.

Questioning the Conventional Approach

Focusing attention on the meeting of the coupling constants (Fig. 1), the question arises: To what extent does this meeting reflect the “truth” or is it somehow deceptive? There are two reasons why such a question is in order.

(1) First, the unity of forces reflected by the meeting of the coupling constants in SUSY SU(5) or SO(10) is truly incomplete, because it comprises only the gauge forces, but not the Higgs-exchange forces. *The latter are still governed by many arbitrary parameters – i.e., the masses, the quartic and the Yukawa couplings of the Higgs bosons – and are thus ununified.* Such arbitrariness goes against the central spirit of grand unification and has been the main reason in my mind since the 1970’s (barring an important caveat due to the growth of superstring theories in the 1980’s, see below) to consider seriously the possibility that the Higgses as well as the quarks and the leptons are composite. Furthermore, neither SUSY SU(5) nor SUSY SO(10), by itself, has the scope of explaining the origins of (a) the three families, (b) inter- and intra-family mass-splittings and (c) the hierarchical mass-scales: from M_{Planck} to m_ν .

(2) The second reason for questioning the conventional approach is this: one might have hoped that one of the two schemes – i.e., the minimal SUSY SU(5) or the SUSY SO(10)-model, or a broken “grand unified” symmetry with relations between its gauge couplings near the string scale, would emerge from one of the solutions of the superstring theories [17,26], which would yield not only the desired spectrum of quarks, leptons and Higgs bosons but also just the right parameters for the Higgs masses as well as their quartic and Yukawa couplings. While it seems

highly nontrivial that so many widely varying parameters should come out in just the right way simply from topological and other constraints of string theories, it would of course be most remarkable if that did happen. *But so far it has not.* There are in fact a very large number of classically allowed degenerate 4D solutions of the superstring theories (Calabi-Yau, orbifold and free fermionic, etc.), although one is not yet able to choose between them. Notwithstanding this general difficulty of a choice, it is interesting that there are at least some three-family solutions. However, not a single one of these has yielded *either* a SUSY SU(5) or an SO(10)-symmetry, *or* a broken “grand unified” symmetry involving direct product of groups, with the desired spectrum *and* Higgs-sector parameters, so as to explain the bizarre pattern of fermion masses and mixings of the three families [27]. Note that for a string theory to yield elementary quarks, leptons and Higgs bosons, either the *entire package* of calculable Higgs-sector parameters, which describe the masses of all the fermions and their mixings (subject to perturbative renormalization), should come out just right, or else the corresponding solution must be discarded. This no doubt is a *heavy burden*. For the case of the broken grand unified models, there is the additional difficulty that the grand unification scale of 2×10^{16} GeV obtained from low-energy extrapolation does not match the string unification scale of about 4×10^{17} GeV [28].

Thus, even if a certain superstring theory is the right starting point, and I believe it is, it is not at all clear, especially in view of the difficulties mentioned above, that it makes contact with the low-energy world by yielding elementary quarks, leptons and Higgs bosons. In this sense, it seems prudent to keep open the possibility that the meeting of the coupling constants in the context of conventional grand unification, which after all corresponds to predicting just one number – i.e., $\sin^2\theta_W$ – correctly, may be fortuitous. Such a meeting should at least be viewed with caution as regards inferring the extent to which it reflects the “truth” because there are in fact alternative ways by which such a meeting can occur (see discussions below).

IV. The Preonic Approach to Unification and Supersymmetry

This brings me to consider an alternative approach to unification based on the ideas of preons and local supersymmetry [9,10,11,12]. Although the general idea of preons is old [30], the particular approach [9-12] which I am about to present has evolved in the last few years. It is still unconventional, despite its promising features. Its lagrangian introduces only six positive and six negative chiral preonic superfields which define the two flavor and four color attributes of a quark-lepton family and possess only *the minimal gauge interactions* corresponding to flavor-color and metacolor gauge symmetries [9]. But the lagrangian *is devoid altogether of the Higgs sector since its superpotential is zero owing to gauge and non-anomalous R-symmetry*. Therefore, it is free from all the arbitrary Higgs-mass, quartic and Yukawa coupling parameters which arise in the conventional approach to grand unification. This brings *real economy*. In fact, the preon model possesses just three (or four) gauge coupling parameters which are the only parameters of the model and even these few would merge into one near the Planck scale if there is an underlying unity of forces as we envisage [29]. By contrast, the standard model has 19 and conventional SUSY grand unification models have over 15 parameters. As mentioned in the introductory chapter, in addition to economy, the main motivations for pursuing the preonic approach are that it provides simple explanations for (a) the protection of the masses of the composite quarks and leptons [10], (b) family replication [11], (c) inter-family mass-hierarchy ($m_{u,d,e} \ll m_{c,s,\mu} \ll m_{t,b,\pi}$) [12], and (d) diverse mass-scales [9]. At the same time, it is viable with respect to observed processes including flavor-changing neutral current processes (see remarks later) and oblique electroweak corrections.

Fermion-boson partnership in a SUSY theory, (i.e. $\psi \leftrightarrow \varphi$ and $v_\mu \leftrightarrow \lambda$ or $\bar{\lambda}$ etc.), leads to several alternative three-particle combinations with identical quantum numbers, which can make a left-chiral $SU(2)_L$ -doublet family q_L^i – e.g. (i) $\sigma_{\mu\nu}\psi_L^f\varphi_R^{c*}v_{\mu\nu}$, (ii) $\sigma^{\mu\nu}\varphi_L^f\psi_R^{c*}v_{\mu\nu}$, (iii) $\psi_L^f\psi_R^{c*}\lambda$ and (iv) $\varphi_L^f(\sigma^\mu\bar{\lambda})\partial_\mu\varphi_R^{c*}$. Here f and c denote flavor and color quantum numbers. *The plurality of these combinations, which stems because of SUSY, is in essence the origin of family-replication*. By constructing composite superfields, Babu, Stremnitzer and I showed [11] that at the level of min-

imum dimensional composite operators (somewhat analogous to qqq for QCD) there are just three linearly independent chiral families $q_{L,R}^i$, and, in addition, two *vector-like families* $Q_{L,R}$ and $Q'_{L,R}$, which couple vectorially to W_L 's and W_R 's respectively. Each of these composite families with spin-1/2 is, of course, accompanied by its scalar superpartner. We thus see that one good answer to Rabi's famous question: "Who ordered that?", is supersymmetry and compositeness.

Certain novel features in the dynamics of a class of SUSY QCD theories, in particular (as mentioned in the introduction) the forbidding of SUSY-breaking in the absence of gravity [8,10], and symmetries of the underlying preonic theory, play crucial roles in obtaining the other desired results – (a), (c) and (d), mentioned above. The reader is referred to the papers in Refs. 9-12 and in particular to a recent review of the preonic approach in Ref. 31 for details of the two broad dynamical assumptions and the reasons underlying a derivation of these results. One attractive feature of the model, which emerges primarily through the symmetries of the underlying lagrangian, is that the two vector-like families $Q_{L,R}$ and $Q'_{L,R}$ (mentioned above) acquire masses of order 1 TeV, while the three chiral families acquire their masses primarily through their spontaneously induced mixings with the two vector-like families. This feature automatically explains why the electron family is so light compared to the tau-family and (owing to additional symmetries) why the masses of the muon-family lie intermediate between those of the electron and the tau-families. In particular, the model explains why $m_e \sim 1 \text{ MeV}$ while $m_t \approx 100 - 180 \text{ GeV}$, i.e., why $(m_e/m_t) \sim 10^{-5}$.

Furthermore, using the values of the standard model gauge couplings measured at LEP and the spectrum of the preon model above and below the preon-binding scale $\Lambda_M \sim 10^{11} \text{ GeV}$, it is found (see Fig. 2) that the flavor-color gauge symmetry being $SU(2)_L \times U(1)_R \times SU(4)^c$ near the Planck scale and the metacolor gauge symmetry being either $SU(5)$ [29] or $SU(6)$ [32], the gauge couplings do tend to meet near the Planck scale. This opens up a novel possibility for grand unification at the preon level and thereby a possible new route for superstring theories to make connection with the low-energy world.

Last but not least the preon model leads to some *crucial predictions* which include the existence of the two vector-like families at the TeV-scale. [See Refs. 12 and 31 for a list of predictions.] These two families can be searched for at the forthcoming LHC, the e^-e^+ next linear collider (in planning) and especially at a future version of the now-extinct SSC. Their discovery or non-discovery with masses up to few TeV will clearly vindicate or exclude the preonic approach developed in Refs. 9-12.

V. Summary and Concluding Remarks

In summary, we see that just by uniting bosons and fermions, Bose-Fermi symmetry ends up in playing an *essential role* in every attempt at higher unification, beyond that of the standard model.

- First, the conventional approach to grand unification, with elementary quarks, leptons and Higgs bosons, of course needs supersymmetry, both for a technical resolution of the gauge hierarchy problem and also to preserve the idea of the meeting of the gauge coupling constants in accord with the LEP data (i.e., precision measurement of $\sin^2 \Theta_W$).

- Second, as we saw the alternative preonic approach to unification requires supersymmetry in a still more crucial manner. The Witten index-theorem, which ensures the protection of supersymmetry and thereby inhibits the formation of supersymmetry and chiral symmetry-breaking condensate $\langle \bar{\psi}\psi \rangle$, in the absence of gravity, plays an essential role in the preonic approach in that it explains why composite quarks and leptons are so light compared to their compositeness scale. In a non-supersymmetric QCD type of theory, there would be no reason for an inhibition of the $\langle \bar{\psi}\psi \rangle$ -condensate. Utilizing local supersymmetry to its advantage, the preonic approach furthermore provides simple explanations for the origins of (i) the three chiral families, (ii) inter-family mass-hierarchy and (iii) diverse mass-scales, and at the same time (iv) provides the scope for a meeting of the gauge coupling constants near the Planck scale. The crucial prediction of the preonic approach – that there

must exist two vector-like families in the TeV-range – is once again tied to supersymmetry. It is because of supersymmetry that both chiral and vector-like families arise as composites naturally from within the model, and also their masses get tied to the supersymmetry-breaking scale.

In short, because supersymmetry provides some novel features in the dynamics relative to non-supersymmetric QCD, such as the protection of chiral symmetry-breaking condensate $\langle \bar{\psi}\psi \rangle$, and the scope for a possible breakdown of parity and global vectorial symmetries like isospin and preon number, it is clear that it would play an even more crucial role in making ends meet if quarks and leptons are composite rather than if they are elementary.

- Last but not least, the idea that the fundamental entities are not point-like but are one-dimensional strings of sizes $\sim 10^{-33}$ cm seems to need Bose-Fermi symmetry first of all to stabilize the vacuum at the string scale, and, second, to avoid large quantum corrections to Higgs (mass)² at long distances. Furthermore, supersymmetry automatically helps avoid tachyons in string theories. Strings combined with supersymmetry give rise to superstrings. As mentioned before, the superstring theories provide the scope for the greatest synthesis so far in particle physics in that they seem capable of unifying *all matter* (spins 0, 1/2, 3/2, 2 and higher) as vibrational modes of the string and also *all their interactions*, which include not only the gauge forces and gravity but also the apparently non-gauge Higgs-type Yukawa and quartic couplings, within a single coherent framework. *The most attractive feature is that the superstring theories permit no dimensionless parameter at the fundamental level.* Equally important, they provide the scope for yielding a good quantum theory of gravity.

For these reasons, I believe that superstring theories possess many (or most) of the crucial ingredients of a “final theory” – “the theory of everything”. But I also believe that, as they stand, they do not constitute the whole of an ultimate theory, because, first and foremost, in spite of the desirable feature that they constrain the gauge symmetry, the spectrum and the S-matrix elements (interactions), they are

not generated by an underlying principle analogous to that of general coordinate or gauge invariance. Second, as a practical matter, they do not yet explain why we live in $3 + 1$ dimensions, and given the fact that supersymmetry does break in the real world, they do not explain why the cosmological constant is so small or zero. Third, they also do not yet provide a consistent understanding of (a) supersymmetry breaking and (b) choice of the ground state. Resolutions of some or all of these latter issues, which may well be inter-related, would clearly involve an understanding of the non-perturbative aspects and the symmetries of superstring dynamics. Recent developments which include the ideas of duality symmetries [33] and the realization that the strong-coupling limit of certain superstring theories is equivalent to the weak-coupling limit of certain other theories [34], permitting the elegant and bold conjecture [35] that there is just one superstring theory, may evolve into a form so as to achieve the lofty goal of solving superstring dynamics. It remains to be seen, however, as to how much of the resolution of the issues mentioned above could come “merely” from our understanding of the non-perturbative dynamics of the existing string theories and how much of such a resolution would involve altogether new ingredients (concepts) at a fundamental level.

As another practical matter, for reasons mentioned in Sections III and IV, it is far from clear that the superstring theories make connections with the low-energy world by yielding elementary quarks, leptons and Higgs bosons. The preonic approach, though unconventional, provides a viable and attractive alternative to the conventional approach. It therefore remains to be seen whether the right superstring theory would yield the elementary quark-lepton-Higgs system with the entire “right package” of Higgs-sector parameters or, instead, the preonic spectrum and the associated gauge symmetry. In the latter case, the superstring theory would, of course, be relieved from yielding the right package of such Higgs sector-parameters because the Higgs-sector is simply absent in the preonic theory.

To conclude, our understanding of superstring theories is rather premature. It would clearly take some time – optimistically a decade but conservatively several

decades –for us to understand (and this may be optimistic) the true nature of superstring theories and to discover the missing ingredients (if any) in these theories, which together would help resolve the issues mentioned above. Meanwhile, regardless of these developments in the future, Bose-Fermi symmetry has clearly evolved as a great synthesizing principle. It is a common denominator and a central feature in all the attempts at higher unification mentioned above. Combined with the idea of strings, it provides the scope, as exhibited in Fig. 3, for unifying matter, forces and mass-scales. *As such, it is hard to imagine how nature could have formulated her laws without the aid of supersymmetry.* It is a concept which, I believe, is here to stay, analogous to those of general coordinate and local gauge invariance. Fortunately, unlike some other concepts, the relevance of Bose-Fermi symmetry to particle physics can be established or falsified, depending upon whether the superpartners are discovered or found to be absent at the forthcoming LEP200, LHC, e^-e^+ NLC and a future version of the now-extinct SSC.

VI. Acknowledgements

The research described in this talk is supported in part by the NSF grant number 9421387. This manuscript was prepared during the author's visit to the Institute for Advanced Study, Princeton, New Jersey, which is supported in part by a sabbatical leave grant from the University of Maryland and in part by a grant by the IAS. It is a pleasure to thank Keith Dienes, Alon Faraggi, and especially Edward Witten for several discussions which were helpful in preparing this manuscript. I wish to thank Partha Ghose and Chanchal Majumdar for arranging an excellent meeting in Calcutta and for their kind hospitality. I would also like to thank Valerie Nowak for her most generous cooperation in typing this manuscript.




Figure 3. A Triple Unification Through Bose-Fermi Symmetry and Strings

Figure 3. A Triple Unification Through Bose-Fermi Symmetry and Strings: Such a unity of forces, families and mass-scales may arise directly through a superstring theory, which could give rise to elementary quarks, leptons and Higgs bosons and fix all the parameters of the standard model at the string unification scale. Alternatively, such a unity may arise through the intermediary of preonic substructures, which may emerge from superstrings. These in turn could generate the three chiral families due to SUSY and a hierarchy of scales owing to inhibition in SUSY-breaking (see text). In either case, the twin ideas of Bose-Fermi symmetry and strings are crucial to achieving this unification.

References

- [1] S.N. Bose, letter to A. Einstein, June 4, 1924; Z. Phys. **26** (1924) 178.
- [2] A. Pais, “Subtle is the Lord,” Oxford Univ. Press, Page 425.
- [3] Initiated (for spin-0 particles) by W. Pauli, letter to W. Heisenberg, June 28, 1934; a review of the different stages of the spin-statistics theorem may be found in R.F. Streater and A.S. Wightman, “PCT, Spin and Statistics and All That,” Benjamin, New York (1964); and in R. Jost in “Theoretical Physics in the Twentieth Century,” eds. M. Fierz and V. Weisskopf, p. 107, Interscience, New York (1960). For a recent review that includes some new developments, see O.W. Greenberg, D.M. Greenberg and T.V. Greenberg in “Quantum Coherence and Reality,” eds. J. Anandan and J.L. Sasko, World Scientific, Singapore (1994), pp. 301-312.
- [4] Y.A. Gelfand and E.S. Likhtman, JETP Lett. **13** (1971) 323; J. Wess and B. Zumino, Nucl. Phys. **B70** (1974)139; Phys. Lett. **49B**(1974) 52; D. Volkov and V.P. Akulov, JETP Lett. **16** (1972) 438.
- [5] R. Haag, J. Lopuszanski and M. Sohnius, Nucl. Phys. **B88** (1975) 61.
- [6] S. Coleman and J. Mandula, Phys. Rev. **159** (1967) 1251.
- [7] E. Witten, Nucl. Phys. **B188** (1981) 513; R. Kaul, Phys. Lett. **109B** (1982) 19; S. Dimopoulos and S. Raby, Nucl. Phys. **B192** (1981) 353.
- [8] E. Witten, Nucl. Phys. **B185** (1981) 513; **B202** (1983) 253; E. Cohen and L. Gomez, Phys. Rev. Lett. **52** (1984) 237.
- [9] J.C. Pati, Phys. Lett. **B228** (1989) 228.
- [10] J.C. Pati, M. Cvetič and H. Sharatchandra, Phys. Rev. Lett. **58** (1987) 851.
- [11] K.S. Babu, J.C. Pati and H. Stremnitzer, Phys. Lett. **B256** (1991) 206.
- [12] K.S. Babu, J.C. Pati and H. Stremnitzer, Phys. Rev. Lett. **67** (1991) 1688.
- [13] J.C. Pati and Abdus Salam; Proc. 15th High Energy Conference, Batavia, reported by J.D. Bjorken, Vol. 2, p. 301 (1972); Phys. Rev. **8** (1973) 1240; Phys. Rev. Lett. **31** (1973) 661; Phys. Rev. **D10** (1974) 275.
- [14] H. Georgi and S.L. Glashow, Phys. Rev. Lett. **32** (1974) 438.

- [15] H. Georgi H. Quinn and S. Weinberg, Phys. Rev. Lett. **33** (1974) 451.
- [16] For a review see E. Farhi and L. Susskind, Phys. Rev. **74** (1981) 277 and references therein.
- [17] M. Green and J. Schwarz, Phys. Lett. **149B** (1984) 117; D. Gross, J. Harvey, E. Martinec and R. Rohm, Phys. Rev. Lett **54** (1985) 502; P. Candelas, G. Horowitz, A. Strominger and E. Witten, Nucl. Phys. **B258** (1985) 46.
- [18] SO(10): H. Georgi, Proc. AIP Conf. Williamsburg (1994); H. Fritzsch and P. Minkowski, Ann. Phys. (NY) **93** (1975) 193. E(6): F. Gursey, P. Ramond and P. Sikivie, Phys. Lett. **60B** (1976) 177.
- [19] Particle Data Group, Review of Particle Properties, Phys. Rev. **D45**, Part II, SI-S584, June 1, 1992.
- [20] LEP data, Particle Data Group (June, 1994).
- [21] S. Dimopoulos and H. Georgi, Nucl. Phys. **B193** (1981) 150; N. Sakai, Z. Phys. **C11** (1982) 153.
- [22] P. Langacker and M. Luo, Phys. Rev. **D44** (1991) 817; U. Amaldi, W. de Boer and H. Furstenau, Phys. Lett. **B260** (1991) 447; J. Ellis, S. Kelley and D.V. Nanopoulos, Phys. Lett. **B260** (1991) 131; F. Anselmo, L. Cifarelli, A. Peterman and A. Zichichi, Nuov. Cim. **A104** (1991) 1817.
- [23] P. Langacker, Review talk at Gatlinburg Conference, June '94, HEP-PH-9411247.
- [24] For analysis of this type, see e.g. R. Arnowitt and P. Nath, Phys. Rev. Lett. **69** (1992) 725; K. Inoue, M. Kawasaki, M. Yamaguchi and T. Yanagida, Phys. Rev. **D45** (1992) 328; G.G. Ross and R.G. Roberts, Nucl. Phys. **B377** (1992) 571; and J.L. Lopez, D. Nanopoulos and H. Pois, Phys. Rev. Lett. **47** (1993) 2468. Other references may be found in the last paper.
- [25] J.C. Pati, A. Salam and U. Sarkar, Phys. Lett. **133B** (1983) 330; J.C. Pati, Phys. Rev. Rap. Comm. **D29** (1983) 1549.
- [26] H. Kawai, D. Lewellen and S. Tye, Nucl. Phys. **B288** (1987) 1; I. Antoniadis, C. Bachas and C. Kounnas, Nucl. Phys. **B289** (1987) 187.

- [27] Some partially successful three-family solutions with top acquiring a mass of the right value ($\approx 175 \text{ GeV}$) and all the other fermions being massless at the level of cubic Yukawa couplings have been obtained with $Z_2 \times Z_2$ orbifold compactification by A. Farragi, Phys. Lett. **B274** (1992) 47; Nucl. Phys. **B416** (1994) 63; and J. Lopez, D. Nanopoulos and A. Zichichi, Texas A&M preprint CTP AMU-06/95). In these attempts, all the other masses and mixings including $m_e \sim \mathcal{O}(1 \text{ MeV})$ are attributed to in-principle calculable higher dimensional operators. It seems optimistic that the entire package of effective parameters would come out correctly this way with the desired hierarchy. But, of course, there is no argument that they cannot. Thus it seems most desirable to pursue this approach as far as one can. This is why I personally keep an open mind with regard to both the conventional approach and the preonic alternative.
- [28] V.S. Kaplunovsky, Nucl. Phys. **B307** (1988) 145. Recently, K.R. Dienes and A.E. Farragi (preprints hep-th/9505018 and hep-th/9505046) provide general arguments why string-threshold corrections arising from the massive tower of states are naturally suppressed and, thus, these corrections do not account for such a mismatch between the two scales. They and other authors have noted that string theories tend to give extra matter, which, if they acquire masses in the right range, could eliminate the mismatch. For a recent discussion of some relevant issues pertaining to string-unification, see these papers as well as L.E. Ibáñez, talk at Strings '95, USC, March 1995, FTUAM 95/15-ReV.
- [29] K.S. Babu and J.C. Pati, Phys. Rev. Rap. Comm. **D48** (1993) R1921.
- [30] Old works on composite models for quarks *and* leptons include the presently-pursued idea of flavon-chromon preons which was introduced in the paper of J.C. Pati and A. Salam, Phys. Rev. **D10** (1974) 275 (Footnote 7). A similar idea that treated only quarks but not leptons as composite was considered independently by O.W. Greenberg (private communication to JCP). This idea has been subsequently considered by Pati and Salam in a set of papers (1975-

80) and by several other authors – with W 's treated as composites in some of them – see e.g., H. Terezawa, Prog. Theor. Phys. **64** (1980) 1763, H. Fritzsch and G. Mandelbaum, Phys. Lett. **102B**, (1981) 113, and O.W. Greenberg and J. Sucher, Phys. Lett. **99B** (1981) 339; and supersymmetric versions in J.C. Pati and A. Salam, Nucl. Phys. **B214** (1983) 109; *ibid.* **B234** (1984) 223 and by R. Barbieri, Phys. Lett. **121B** (1983) 43. None of these works provided a reason, however, for (a) the protection of composite quark-lepton masses, (b) family-replication and (c) inter-family mass-hierarchy. The interesting idea of quasi-Nambu Goldstone fermions suggested by W. Büchmüller, R. Peccei and T. Yanagida, Phys. Lett. **124B** (1983) 67, provided a partial reason for some of these issues, in particular for (a), but had problems of internal consistency as regards SUSY-breaking while maintaining the lightness of composite quarks and generating effective gauge symmetries. The idea of SUSY-compositeness developed in Refs. 9-12 and Ref. 29 introduces a *new phase* in the preonic approach in that (i) it avoids the problems of technicolor and (ii) it seems capable of incorporating the idea of grand unification [29], while providing a reason for each of the issues (a), (b) and (c) mentioned above. It is these features which seem to make the new approach a *viable alternative* to the conventional approach to grand unification.

- [31] For a recent review of the preonic approach, see J.C. Pati “Towards a Unified Origin of Forces, Families and Mass Scales,” hep-ph/9505227, to appear in the Proc. of the 1994 Int’l. Conf. on B-Physics, held at Nagoya, Japan (Oct. 26-29, 1994).
- [32] The threshold effects at Λ_M which permit unity with metacolor symmetry being $SU(6)$ have been considered by K.S. Babu, J.C. Pati and M. Parida (to appear).
- [33] See e.g., A. Sen, Int. J. Mod. Phys. **A9** (1994) 3707, hep-th/9402002 and hep-th/9402032; J.H. Schwarz, hep-th/9411178; C.M. Hull and P.K. Townsend, QMW94-39, R/94/33 and references therein.

- [34] E. Witten, preprint IASSNS-HEP-95/18, hep-th/9503124.
- [35] E. Witten, Ref. 34 and private communications.